

Kearon C, Akl EA, Comerota AJ, Prandoni P, Bounameaux H, Goldhaber SZ, Nelson ME, Wells PS, Gould MK, Dentall F, Crowther M, Kahn SR: American College of Chest Physicians. Antithrombotic therapy for VTE disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines.[Erratum appears in Chest. 2012 Dec;142(6):1698-1704]. Chest. 2012 Feb;141(2 Suppl):e419S–94S.

Chalmers E, Ganesan V, Liesner R, Maroo S, Nokes T, Saunders D, Williams M: British Committee for Standards in Haematology. Guideline on the investigation, management and prevention of venous thrombosis in children. Br J Haematol. 2011 Jul;154(2):196–207.

Monagle P, Chan AK, Goldenberg NA, Ichord RN, Journeycake JM, Nowak-Göttl U, Vesely SK: American College of Chest Physicians. Antithrombotic therapy in neonates and children: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest. 2012 Feb;141(2 Suppl):e737S–801S.

Evidence-based management of sickle cell disease: expert panel report, 2014. Washington (DC): National Institutes of Health, National Heart, Lung, and Blood Institute; 2014. 161 p.

Blood transfusion guideline. Dutch Institute for Healthcare Improvement CBO; 2011. 402 p.

Oscier D, Dearden C, Eren E, Fegan C, Follows G, Hillmen P, Illidge T, Malutes E, Milligan DW, Pettitt A, Schuh A, Wimperis J: British Committee for Standards in Haematology. Guidelines on the diagnosis, investigation and management of chronic lymphocytic leukaemia. Br J Haematol. 2012 Dec;159(5):541–64.

Eichhorst B, Hallek M, Dreyling M, Group EGW. Chronic lymphocytic leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol. 2010 May;21 Suppl 5:v162–4.

Watson H, Davidson S, Keeling D. Guidelines on the diagnosis and management of heparin-induced thrombocytopenia: second edition. Br J Haematol. 2012;159(5):528–40.

Cuker A, Gimotty PA, Crowther MA, Warkentin TE. Predictive value of the 4Ts scoring system for heparin-induced thrombocytopenia: a systematic review and meta-analysis. Blood. 2012;120:4160–7.

Don't transfuse more than the minimum number of red blood cell (RBC) units necessary to relieve symptoms of anemia or to return a patient to a safe hemoglobin range (7 to 8 g/dL in stable, non-cardiac in-patients).

Transfusion of the smallest effective dose of RBCs is recommended because liberal transfusion strategies do not improve outcomes when compared to restrictive strategies. Unnecessary transfusion generates costs and exposes patients to potential adverse effects without any likelihood of benefit. Clinicians are urged to avoid the routine administration of 2 units of RBCs if 1 unit is sufficient and to use appropriate weight-based dosing of RBCs in children.

Don't test for thrombophilia in adult patients with venous thromboembolism (VTE) occurring in the setting of major transient risk factors (surgery, trauma or prolonged immobility).

Thrombophilia testing is costly and can result in harm to patients if the duration of anticoagulation is inappropriately prolonged for patients who are incorrectly labeled as thrombophilic. Thrombophilia testing does not change the management of VTEs occurring in